


AHPWA24AUSA

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1(Original). A method of inducing contraception comprising the step of delivering to a female of child-bearing age a composition comprising a compound of formula I, or a tautomer thereof, in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said female, wherein formula I is:



I

wherein:

R^1 and R^2 are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

AHPWA24AUSA

p is an integer from 1 to 4;

or R¹ and R² form a double bond to C(CH₃)₂, C(cycloalkyl), O, or C(cycloether);


R³ is selected from the group consisting of H, OH, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₆ alkenyl, substituted C₃ to C₆ alkenyl, alkynyl, substituted alkynyl, and COR^A;

R^A is selected from the group consisting of H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, and substituted C₁ to C₃ aminoalkyl;

R⁴ is selected from the group consisting of H, halogen, CN, NH₂, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₁ to C₆ alkoxy, substituted C₁ to C₆ alkoxy, C₁ to C₆ aminoalkyl, and substituted C₁ to C₆ aminoalkyl;

R⁵ is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:



X is selected from the group consisting of halogen, OH, CN, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ thioalkyl, substituted C₁ to C₃ thioalkyl, S(O)alkyl, S(O)₂alkyl, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, NO₂, C₁ to C₃ perfluoroalkyl, substituted C₁ to C₃ perfluoroalkyl, 5 or 6 membered heterocyclic ring having 1 to 3 heteroatoms, CONH₂, CSNH₂, CNHNHOH, CNH₂NOH, CNHNOH, COR^B, CSR^B, OCOR^B, and NR^CCOR^B;

R^B is selected from the group consisting of H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, and substituted C₁ to C₃ aminoalkyl;

R^C is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

Y and Z are independently selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ thioalkyl, and substituted C₁ to C₃ thioalkyl;

AHPWA24AUSA

b) a five or six membered heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR⁶ and having one or two independent substituents from the group consisting of H, halogen, CN, NO₂, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, COR^D, CSR^D, and NR^ECOR^D;

R^D is H, NH₂, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R⁶ is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, or C₁ to C₄CO₂alkyl; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group consisting of halogen, alkyl, substituted alkyl, CN, NO₂, alkoxy, substituted alkoxy, and CF₃;


Q¹ is S, NR⁷, or CR⁸R⁹;

R⁷ is selected from the group consisting of CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO₂CF₃, OR¹¹, and NR¹¹R¹²;

R⁸ and R⁹ are independent substituents selected from the group consisting of H, alkyl, substituted alkyl, acyl, substituted acyl, aroyl, substituted aroyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO₂, CN, and CO₂R¹⁰;

R¹⁰ is C₁ to C₃ alkyl or substituted C₁ to C₃ alkyl;

or CR⁸R⁹ comprise a six membered ring having the structure:



AHPWA24AUSA

R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

2(Original). The method according to claim 1, wherein said compound of formula I and said selective estrogen receptor modulator are delivered in a single composition.

3(Original). The method according to claim 1, wherein said compound of formula I and said selective estrogen receptor modulator are delivered separately.

4(Original). The method according to claim 1, wherein said selective estrogen receptor modulator is selected from the group consisting of EM-800, EM-652, raloxifene hydrochloride, arzoxifene, lasofoxifene, droloxifene, tamoxifen citrate, 4-hydroxytamoxifen citrate, clomiphene citrate, toremifene citrate, pipendoxifene, idoxifene, levormeloxifene, centchroman, nafoxidene, and bazedoxifene.


5(Original). The method according to claim 1, wherein said compound is delivered at a daily dosage of about 0.1 to about 50 mg.

6(Original). The method according to claim 1, wherein said regimen comprises delivering said composition daily for 1 to about 21 days, wherein said regimen is a cycle which is repeated monthly.

7(Original). The method according to claim 1, wherein said selective estrogen receptor modulator is delivered at a daily dosage of about 0.2 to about 100 mg.

AHPWA24AUSA

8(Original). The method according to Claim 1, wherein R^1 and R^2 are joined to form a $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$ ring; n is 3; R^3 and R^4 are H; R^5 is the substituted benzene ring having the structure:




X is selected from the group consisting of halogen, CN, CONH_2 , CSNH_2 , COR^B , CSR^B , C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring having 1 to 3 heteroatoms, and C_1 to C_3 thioalkyl;

R^B is C_1 to C_3 aminoalkyl or substituted C_1 to C_3 aminoalkyl, wherein said aminoalkyl is $\text{NH}(\text{alkyl})$ or $\text{N}(\text{alkyl})_2$;

Y is selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, C_1 to C_4 alkyl, and C_1 to C_3 thioalkyl.

9(Original). The method according to Claim 1, wherein R^1 and R^2 are joined to form the $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$ ring; n is 3; R^3 and R^4 are H; R^5 is the five membered ring having the structure:



U is O, S, or NR^6 ;


X' is selected from the group consisting of halogen, CN, NO_2 , CONH_2 , CSNH_2 , COR^B , CSR^B , C_1 to C_3 alkyl, and C_1 to C_3 alkoxy;

R^B is C_1 to C_3 aminoalkyl or substituted C_1 to C_3 aminoalkyl, wherein said aminoalkyl is $\text{NH}(\text{alkyl})$ or $\text{N}(\text{alkyl})_2$;

Y' is selected from the group consisting of H, halogen, and C_1 to C_4 alkyl, wherein said halogen is F.

AHPWA24AUSA

10(Original). The method according to Claim 1, wherein R^1 and R^2 are joined to form a $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$ ring; n is 3; R^3 and R^4 are H; R^5 is the six membered ring having the structure:



X^1 is N or CX^2 ;

X^2 is halogen, CN, CONH_2 , CSNH_2 , COR^B , CSR^B , or NO_2 ;

R^B is C_1 to C_3 aminoalkyl or substituted C_1 to C_3 aminoalkyl, wherein said aminoalkyl is $\text{NH}(\text{alkyl})$ or $\text{N}(\text{alkyl})_2$.

11(Original). The method according to claim 1, wherein R^1 and R^2 are alkyl or substituted alkyl; R^3 is H.

12(Original). The method according to claim 1, wherein R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$; R^3 is H.

13(Original). The method according to claim 1, wherein R^3 is H; Q^1 is S or NR^7 .

14(Original). The method according to claim 1, wherein said compound is selected from the group consisting of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1H)-thione, 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzonitrile, 4-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3'-[3H]indol]-5-yl)-5-fluorobenzonitrile, 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3'-[3H]indol]-5-yl)-2-thiophenethioamide, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-

AHPWA24AUSA

[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-H-pyrrole-2-carbonitrile, 5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3-thiophenecarbonitrile, 5-(1,2-Dihydro-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-2-thiophenecarbonitrile, 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-Benzyl-5-(3-chlorophenyl)-3-methyl-1,3-dihydro-2H-indole-2-thione, 4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile, 5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile, 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile, 5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile, 5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile, 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-4-methyl-2-thiophenecarbonitrile, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-2-thiophenecarbonitrile, 5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-

AHPWA24AUSA


chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione, 5-(4-Fluoro-3-(trifluoromethyl)phenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-n-butyl-2-thiophenecarbonitrile, 5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine, N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine, 5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(2-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(4-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 3-(1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile, 5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(acetoxylimino)-5'-yl)-2-thiophenecarbonitrile, 3-Fluoro-N'-hydroxy-5-(2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide, N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indol]-2'-hydroxyimino)-5'-yl-2-thiophenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboximidamide, 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-1H-

AHPWA24AUSA

pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2-ylidenecyanamide, 5'-(5-Cyano-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-methyl-thiophene-2-carbonitrile, and 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

15(Currently Amended). The method according to ~~any of claims~~ claim 1, wherein said compound is 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

16(Original). A method of inducing contraception comprising the step of delivering to a female of child-bearing age a composition comprising a compound of formula II, or a tautomer thereof, in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said female, wherein formula II is:




wherein:

AHPWA24AUSA

R^{11} is selected from the group consisting of H, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl;

R^5 is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:




wherein:

X is selected from the group consisting of halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNHOH, CNH₂NOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring having 1 to 3 heteroatoms, and C₁ to C₃ thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkyl;

(ii) a five membered ring having the structure:



wherein:

U is O, S, or NR⁶;


R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CNHNHOH, CNH₂NOH, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F, and C₁ to C₄ alkyl; or

(iii) a six membered ring having the structure:

AHPWA24AUSA



wherein:


X^1 is N or CX^2 ;

X^2 is halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CON(alkyl)_2$, $CSN(alkyl)_2$ or NO_2 ;

or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

17(Original). The method according to claim 16, wherein R^5 is said five membered ring and U is O or S.


18(Original). A method of inducing contraception comprising the step of delivering to a female of child-bearing age a composition comprising a compound of formula III in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said female, wherein formula III is:



wherein:

R^5 is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:



AHPWA24AUSA

wherein:

X is selected from the group consisting of halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring having 1 to 3 heteroatoms, and C₁ to C₃ thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkyl;

(ii) a five membered ring having the structure:



wherein:


U is O, S, or NR⁶;

R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F and C₁ to C₄ alkyl; or

(iii) a six membered ring having the structure:



wherein:

X¹ is N or CX²;


X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂ or NO₂;

or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

AHPWA24AUSA

19(Original). The method according to claim 17, wherein R^5 is the five membered ring (ii) and U is O or S.

20(Original). A method of inducing contraception comprising the step of delivering to a female of child-bearing age a composition comprising a compound of formula IV in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said female, wherein formula IV is:




wherein:

R^8 is selected from the group consisting of H, CO_2R^{10} , acyl, substituted acyl, aroyl, substituted aroyl, alkyl, substituted alkyl, and CN;

R^{10} is C_1 to C_3 alkyl;

R^5 is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:



wherein:

X is selected from the group consisting of halogen, CN, $CONH_2$, $CSNH_2$, $CONHalkyl$, $CSNHalkyl$, $CON(alkyl)_2$, $CSN(alkyl)_2$, $CNHNOH$, C_1 to C_3 alkoxy, C_1 to C_3 alkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 membered heterocyclic ring having 1 to 3 heteroatoms, and C_1 to C_3 thioalkyl;

AHPWA24AUSA

Y is selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkyl;

(ii) a five membered ring having the structure:



wherein:


U is O, S, or NR⁶;

R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;

X' is selected from the group consisting of halogen, CN, NO₂, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F and C₁ to C₄ alkyl;

(iii) a six membered ring having the structure:



wherein:

X¹ is N or CX²;

X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂ or NO₂;


or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

21(Original). The method according to claim 10, wherein R⁵ is the five-membered ring (ii) and U is O or S.

22(Original). A method of inducing contraception comprising the step of delivering to a female of child-bearing age a composition comprising a compound of formula V in a regimen which involves delivering a pharmaceutically effective amount of


AHPWA24AUSA

one or more of a selective estrogen receptor modulator to said female, wherein formula V is:



R^5 is (i), (ii), or (iii):

(i) a substituted benzene ring having the structure:




wherein:

X is selected from the group consisting of halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, CNHNOH, C₁ to C₃ alkoxy, C₁ to C₃ alkyl, NO₂, C₁ to C₃ perfluoroalkyl, 5 membered heterocyclic ring having 1 to 3 heteroatoms, and C₁ to C₃ thioalkyl;

Y is selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, C₁ to C₄ alkyl, and C₁ to C₃ thioalkyl;

(ii) a five membered ring having the structure:



wherein:

U is O, S, or NR⁶;

R⁶ is H, C₁ to C₃ alkyl, or C₁ to C₄ CO₂alkyl;


X' is selected from the group consisting of halogen, CN, NO₂, CONH₂,

AHPWA24AUSA

CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂, C₁ to C₃ alkyl, and C₁ to C₃ alkoxy;

Y' is selected from the group consisting of H, F, and C₁ to C₄ alkyl;

(iii) a six membered ring having the structure:



wherein:


X¹ is N or CX²;

X² is halogen, CN, CONH₂, CSNH₂, CONHalkyl, CSNHalkyl, CON(alkyl)₂, CSN(alkyl)₂ or NO₂;

or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

23(Original). The method according to claim 22, wherein R⁵ is the five membered ring (ii) and U is O or S.

24(Withdrawn). A method of providing hormone replacement therapy comprising the step of delivering to a female a composition comprising a compound of formula I, or a tautomer thereof, in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said female, wherein formula I is:



wherein:

R¹ and R² are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring,

AHPWA24AUSA

substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R^1 and R^2 form a double bond to $\text{C}(\text{CH}_3)_2$, $\text{C}(\text{cycloalkyl})$, O , or $\text{C}(\text{cycloether})$;


R^3 is selected from the group consisting of H , OH , NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_6 alkenyl, substituted C_3 to C_6 alkenyl, alkynyl, substituted alkynyl, and COR^A ;

R^A is selected from the group consisting of H , C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^4 is selected from the group consisting of H , halogen, CN , NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_1 to C_6 alkoxy, substituted C_1 to C_6 alkoxy, C_1 to C_6 aminoalkyl, and substituted C_1 to C_6 aminoalkyl;

R^5 is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:



X is selected from the group consisting of halogen, OH , CN , C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 thioalkyl, substituted C_1 to C_3 thioalkyl, $\text{S}(\text{O})\text{alkyl}$, $\text{S}(\text{O})_2\text{alkyl}$, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, NO_2 , C_1 to C_3 perfluoroalkyl, substituted C_1 to C_3

AHPWA24AUSA

perfluoroalkyl, 5 or 6 membered heterocyclic ring having 1 to 3 heteroatoms, CONH_2 , CSNH_2 , CNHNHOH , CNH_2NOH , CNHNOH , COR^B , CSR^B , OCOR^B , and NR^CCOR^B ;

R^B is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^C is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

Y and Z are independently selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_4 alkyl, substituted C_1 to C_4 alkyl, C_1 to C_3 thioalkyl, and substituted C_1 to C_3 thioalkyl;

b) a five or six membered heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO_2 and NR^6 and having one or two independent substituents from the group consisting of H, halogen, CN, NO_2 , C_1 to C_4 alkyl, substituted C_1 to C_4 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, COR^D , CSR^D , and NR^ECOR^D ;

R^D is H, NH_2 , C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, or substituted C_1 to C_3 aminoalkyl;

R^E is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

R^6 is H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, or C_1 to $\text{C}_4\text{CO}_2\text{alkyl}$; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group consisting of halogen, alkyl, substituted alkyl, CN, NO_2 , alkoxy, substituted alkoxy, and CF_3 ;

Q^1 is S, NR^7 , or CR^8R^9 ;


R^7 is selected from the group consisting of CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO_2CF_3 , OR^{11} , and $\text{NR}^{11}\text{R}^{12}$;

AHPWA24AUSA

R^8 and R^9 are independent substituents selected from the group consisting of H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO_2 , CN, and CO_2R^{10} ;

R^{10} is C_1 to C_3 alkyl or substituted C_1 to C_3 alkyl;

or CR^8R^9 comprise a six membered ring having the structure:



R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

25(Withdrawn). The method according to claim 23, wherein said hormone replacement therapy is perimenopausal, menopausal, or postmenopausal.

26(Withdrawn). The method according to claim 24, wherein R^1 and R^2 are alkyl or substituted alkyl; R^3 is H.

27(Withdrawn). The method according to claim 24, wherein R^1 and R^2 are joined to form a ring selected from the group consisting of $-CH_2(CH_2)_nCH_2-$, $-CH_2CH_2C(CH_3)_2CH_2CH_2-$, $-O(CH_2)_mCH_2-$, $-O(CH_2)_pO-$, $-CH_2CH_2OCH_2CH_2-$, $-CH_2CH_2N(H)CH_2CH_2-$, and $-CH_2CH_2N(alkyl)CH_2CH_2-$; R^3 is H.

28(Withdrawn). The method according to claim 24, wherein R^3 is H; Q^1 is S or NR^7 .

AHPWA24AUSA

29(Withdrawn). The method according to claim 24, wherein said compound is selected from the group consisting of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1H)-thione, 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzonitrile, 4-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophenethioamide, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-H-pyrrole-2-carbonitrile, 5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3-thiophenecarbonitrile, 5-(1,2-Dihydro-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-2-thiophenecarbonitrile, 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-Benzyl-5-(3-chlorophenyl)-3-methyl-1,3-dihydro-2H-indole-2-thione, 4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile, 5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile, 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile, 5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile, 5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-

AHPWA24AUSA

thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile, 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-4-methyl-2-thiophenecarbonitrile, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-2-thiophenecarbonitrile, 5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione, 5-(4-Fluoro-3-(trifluoromethyl)phenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-n-butyl-2-thiophenecarbonitrile, 5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3''-[3H]indol]-2-amine, N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine, 5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(2-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(4-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 3-(1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile,


AHPWA24AUSA

5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(acetoxyimino)-5'-yl)-2-thiophenecarbonitrile, 3-Fluoro-N'-hydroxy-5-(2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide, N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indol]-2'-hydroxyimino)-5'-yl-2-thiophenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboximidamide, 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-methyl-thiophene-2-carbonitrile, and 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

30(Withdrawn). The method according to claim 24, wherein said compound is 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

AHPWA24AUSA

31(Withdrawn). A method of treating carcinomas comprising the step of delivering to a mammal in need thereof a composition comprising a compound of formula I, or a tautomer thereof, in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said mammal, wherein formula I is:



wherein:

R^1 and R^2 are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R^1 and R^2 form a double bond to $\text{C}(\text{CH}_3)_2$, $\text{C}(\text{cycloalkyl})$, O, or $\text{C}(\text{cycloether})$;

R^3 is selected from the group consisting of H, OH, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_6 alkenyl, substituted C_3 to C_6 alkenyl, alkynyl, substituted alkynyl, and COR^A ;


R^A is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

AHPWA24AUSA

R^4 is selected from the group consisting of H, halogen, CN, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_1 to C_6 alkoxy, substituted C_1 to C_6 alkoxy, C_1 to C_6 aminoalkyl, and substituted C_1 to C_6 aminoalkyl;

R^5 is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:



X is selected from the group consisting of halogen, OH, CN, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 thioalkyl, substituted C_1 to C_3 thioalkyl, $S(O)alkyl$, $S(O)_2alkyl$, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, NO_2 , C_1 to C_3 perfluoroalkyl, substituted C_1 to C_3 perfluoroalkyl, 5 or 6 membered heterocyclic ring having 1 to 3 heteroatoms, $CONH_2$, $CSNH_2$, $CNHNHOH$, CNH_2NOH , $CNHNOH$, COR^B , CSR^B , $OCOR^B$, and $NR^C COR^B$;

R^B is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^C is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

Y and Z are independently selected from the group consisting of H, halogen, CN, NO_2 , C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_4 alkyl, substituted C_1 to C_4 alkyl, C_1 to C_3 thioalkyl, and substituted C_1 to C_3 thioalkyl;

b) a five or six membered heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO_2 and NR^6 and having one or two independent substituents from the group consisting of H, halogen, CN, NO_2 , C_1 to C_4 alkyl, substituted C_1 to C_4 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, COR^D , CSR^D , and $NR^E COR^D$;

R^D is H, NH_2 , C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, or substituted C_1 to C_3 aminoalkyl;

AHPWA24AUSA

R^E is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

R^6 is H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, or C_1 to C_4CO_2 alkyl; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group consisting of halogen, alkyl, substituted alkyl, CN, NO_2 , alkoxy, substituted alkoxy, and CF_3 ;


Q^1 is S, NR^7 , or CR^8R^9 ;

R^7 is selected from the group consisting of CN, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO_2CF_3 , OR^{11} , and $NR^{11}R^{12}$;

R^8 and R^9 are independent substituents selected from the group consisting of H, C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_8 cycloalkyl, substituted C_3 to C_8 cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO_2 , CN, and CO_2R^{10} ;

R^{10} is C_1 to C_3 alkyl or substituted C_1 to C_3 alkyl;

or CR^8R^9 comprise a six membered ring having the structure:



R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

32(Withdrawn). The method according to claim 31, wherein said carcinomas are selected from the group consisting of ovary, breast, colon, endometrial, uterine, and prostate carcinomas.

AHPWA24AUSA

33(Withdrawn). The method according to claim 31, wherein R^1 and R^2 are alkyl or substituted alkyl; R^3 is H.

34(Withdrawn). The method according to claim 31, wherein R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$; R^3 is H.

35(Withdrawn). The method according to claim 31, wherein R^3 is H; Q^1 is S or NR^7 .

36(Withdrawn). The method according to claim 31, wherein said compound is selected from the group consisting of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1H)-thione, 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzonitrile, 4-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophenethioamide, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-H-pyrrole-2-carbonitrile, 5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3-thiophenecarbonitrile, 5-(1,2-Dihydro-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-2-thiophenecarbonitrile, 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-Benzyl-5-(3-chlorophenyl)-3-methyl-

AHPWA24AUSA

1,3-dihydro-2H-indole-2-thione, 4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile, 5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile, 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile, 5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile, 5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile, 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-4-methyl-2-thiophenecarbonitrile, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-2-thiophenecarbonitrile, 5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione, 5-(4-Fluoro-3-(trifluoromethyl)phenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-n-butyl-2-thiophenecarbonitrile, 5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine, N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine, 5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(2-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(4-

AHPWA24AUSA


Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 3-(1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile, 5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'(acetoxymino)-5'-yl)-2-thiophenecarbonitrile, 3-Fluoro-N'-hydroxy-5-(2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide, N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboximidamide, 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-

AHPWA24AUSA

methyl-thiophene-2-carbonitrile, and 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

37(Withdrawn). The method according to claim 31, wherein said compound is 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

38(Withdrawn). A method of treating dysfunctional bleeding, uterine leiomyomata, endometriosis, and/or polycystic ovary syndrome, comprising the step of delivering to a female in need thereof a composition comprising a compound of formula I, or a tautomer thereof, in a regimen which involves delivering a pharmaceutically effective amount of one or more of a selective estrogen receptor modulator to said female, wherein formula I is:



I

wherein:

R^1 and R^2 are selected from the group consisting of H, alkyl, substituted alkyl, OH, O(alkyl), O(substituted alkyl), O(Acetyl), aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, alkylaryl, substituted alkylaryl, alkylheteroaryl, substituted alkylheteroaryl, 1-propynyl, substituted 1-propynyl, 3-propynyl, and substituted 3-propynyl;

or R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$;

AHPWA24AUSA

m is an integer from 1 to 4;

n is an integer from 1 to 5;

p is an integer from 1 to 4;

or R^1 and R^2 form a double bond to $C(CH_3)_2$, $C(\text{cycloalkyl})$, O, or $C(\text{cycloether})$;


R^3 is selected from the group consisting of H, OH, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_3 to C_6 alkenyl, substituted C_3 to C_6 alkenyl, alkynyl, substituted alkynyl, and COR^A ;

R^A is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^4 is selected from the group consisting of H, halogen, CN, NH_2 , C_1 to C_6 alkyl, substituted C_1 to C_6 alkyl, C_1 to C_6 alkoxy, substituted C_1 to C_6 alkoxy, C_1 to C_6 aminoalkyl, and substituted C_1 to C_6 aminoalkyl;

R^5 is selected from the group consisting of a), b) and c):

a) a substituted benzene ring having the structure:



X is selected from the group consisting of halogen, OH, CN, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 thioalkyl, substituted C_1 to C_3 thioalkyl, $S(O)alkyl$, $S(O)_2alkyl$, C_1 to C_3 aminoalkyl, substituted C_1 to C_3 aminoalkyl, NO_2 , C_1 to C_3 perfluoroalkyl, 5 or 6 membered heterocyclic ring having 1 to 3 heteroatoms, $CONH_2$, $CSNH_2$, $CNHNHOH$, CNH_2NOH , $CNHNHOH$, COR^B , CSR^B , $OCOR^B$, and $NR^C COR^B$;

R^B is selected from the group consisting of H, C_1 to C_3 alkyl, substituted C_1 to C_3 alkyl, aryl, substituted aryl, C_1 to C_3 alkoxy, substituted C_1 to C_3 alkoxy, C_1 to C_3 aminoalkyl, and substituted C_1 to C_3 aminoalkyl;

R^C is H, C_1 to C_3 alkyl, or substituted C_1 to C_3 alkyl;

AHPWA24AUSA

Y and Z are independently selected from the group consisting of H, halogen, CN, NO₂, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ thioalkyl, and substituted C₁ to C₃ thioalkyl;

b) a five or six membered heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of O, S, SO, SO₂ and NR⁶ and having one or two independent substituents from the group consisting of H, halogen, CN, NO₂, C₁ to C₄ alkyl, substituted C₁ to C₄ alkyl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, substituted C₁ to C₃ aminoalkyl, COR^D, CSR^D, and NR^ECOR^D;

R^D is H, NH₂, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, aryl, substituted aryl, C₁ to C₃ alkoxy, substituted C₁ to C₃ alkoxy, C₁ to C₃ aminoalkyl, or substituted C₁ to C₃ aminoalkyl;

R^E is H, C₁ to C₃ alkyl, or substituted C₁ to C₃ alkyl;

R⁶ is H, C₁ to C₃ alkyl, substituted C₁ to C₃ alkyl, or C₁ to C₄CO₂alkyl; or

c) an indol-4-yl, indol-7-yl or benzo-2-thiophene moiety, wherein said moiety is optionally substituted by from 1 to 3 substituents selected from the group consisting of halogen, alkyl, substituted alkyl, CN, NO₂, alkoxy, substituted alkoxy, and CF₃;

Q¹ is S, NR⁷, or CR⁸R⁹;


R⁷ is selected from the group consisting of CN, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, SO₂CF₃, OR¹¹, and NR¹¹R¹²;

R⁸ and R⁹ are independent substituents selected from the group consisting of H, C₁ to C₆ alkyl, substituted C₁ to C₆ alkyl, C₃ to C₈ cycloalkyl, substituted C₃ to C₈ cycloalkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, NO₂, CN, and CO₂R¹⁰;

R¹⁰ is C₁ to C₃ alkyl or substituted C₁ to C₃ alkyl;

or CR⁸R⁹ comprise a six membered ring having the structure:

AHPWA24AUSA



R^{11} and R^{12} are independently selected from the group consisting of H, alkyl, substituted alkyl, aryl, substituted aryl, heterocyclic ring, substituted heterocyclic ring, acyl, substituted acyl, aroyl, substituted aroyl, sulfonyl, and substituted sulfonyl; or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

39(Withdrawn). The method according to claim 38, wherein R^1 and R^2 are alkyl or substituted alkyl; R^3 is H.

40(Withdrawn). The method according to claim 38, wherein R^1 and R^2 are joined to form a ring selected from the group consisting of $-\text{CH}_2(\text{CH}_2)_n\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2-$, $-\text{O}(\text{CH}_2)_m\text{CH}_2-$, $-\text{O}(\text{CH}_2)_p\text{O}-$, $-\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{N}(\text{H})\text{CH}_2\text{CH}_2-$, and $-\text{CH}_2\text{CH}_2\text{N}(\text{alkyl})\text{CH}_2\text{CH}_2-$; R^3 is H.

41(Withdrawn). The method according to claim 38, wherein R^3 is H; Q^1 is S or NR^7 .

42(Withdrawn). The method according to claim 38, wherein said compound is selected from the group consisting of 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-thione, 3-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzonitrile, 4-(1',2'-Dihydro-2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-2-thiophenecarbonitrile, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 4-Methyl-5-(1,2-dihydro-2-thioxospiro[cyclohexane-1,3-[3H]-indol]-5-yl)-2-thiophenethioamide, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-(tert-butoxycarbonyl)-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-

AHPWA24AUSA

thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-1-H-pyrrole-2-carbonitrile, 5-(2'-thioxospiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-pyrrole-2-carbonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-3-thiophenecarbonitrile, 5-(1,2-Dihydro-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-2-thiophenecarbonitrile, 5-(3-Fluoro-4-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(2-Amino-5-pyrimidinyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclopentane-1,3-[3H]indol]-5-yl)-5-fluorobenzonitrile, 5-(3-chlorophenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-Benzyl-5-(3-chlorophenyl)-3-methyl-1,3-dihydro-2H-indole-2-thione, 4-(3,3-dimethyl-2-thioxo-2,3-dihydro-1H-indol-5-yl)-2-furonitrile, 5-(3-methoxyphenyl)-3,3-dimethyl-1,3-dihydro-2H-indole-2-thione, 3-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-pyridinecarbonitrile, 5-(3,4-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(5-Chloro-2-thienyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-3-furancarbonitrile, 5-(3-Chloro-4-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chloro-5-fluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3,5-Difluorophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-propyl-2-thiophenecarbonitrile, 5-(3-Fluoro-4-nitrophenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-furancarbonitrile, 5''-(3-Chlorophenyl)spiro[cyclobutane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(2-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5''-(4-Chlorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-4-methyl-2-thiophenecarbonitrile, 5-(1'',2''-Dihydro-2''-thioxospiro[cyclohexane-1,3''-[3H]indol]-5''-yl)-2-thiophenecarbonitrile, 5''-(3-Fluorophenyl)spiro[cyclohexane-1,3''-[3H]indol]-2''(1''H)-thione, 5-(3-Hydroxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-chlorophenyl)-3,3-diethyl-1,3-dihydro-2H-indole-2-thione, 5-(4-Fluoro-3-

AHPWA24AUSA

(trifluoromethyl)phenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 4-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-2-fluorobenzonitrile, 5-(1,2-Dihydro-2-thioxospiro[cyclohexane-1,3-[3H]indol]-5-yl)-4-n-butyl-2-thiophenecarbonitrile, 5-(3-Fluoro-5-methoxyphenyl)spiro[cyclohexane-1,3-[3H]indol]-2(1H)-thione, 5-(3-Chlorophenyl)-N-hydroxyspiro[cyclohexane-1,3'-[3H]indol]-2-amine, N-(Acetyloxy)-5'-(3-chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2''amine, 5'-(3-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(2-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(4-Fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3,4-difluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-methoxyphenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-nitrophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 5'-(3-cyanophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'(1'H)-one oxime, 3-(1',2'-Dihydro-2'-(hydroxyimino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluorobenzonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarbonitrile, 5-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-1-methyl-2-carbonitrile, 5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-1H-pyrrole-2-carbonitrile, 4-(Spiro[cyclohexane-1,3'-[3H]indol]-2'-(acetoxymino)-5'-yl)-2-thiophenecarbonitrile, 3-Fluoro-N'-hydroxy-5-(2'-(hydroxyamino)spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)benzenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-4-methyl-2-thiophenecarboximidamide, N'-Hydroxy-4-(spiro[cyclohexane-1,3'-[3H]indol]-2'-hydroxyimino)-5'-yl-2-thiophenecarboximidamide, N'-Hydroxy-5-(spiro[cyclohexane-1,3'-[3H]indol]-2'-(hydroxyimino)-5'-yl)-2-thiophenecarboxidamide, 5'-(3-Chlorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(3-Cyano-5-fluorophenyl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-

AHPWA24AUSA

thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-3-methyl-thiophen-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 5'-(5-Cyano-thiophen-3-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, 3-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-5-fluoro-benzonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-1-methyl-1H-pyrrole-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, 5-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-4-methyl-thiophene-2-carbonitrile, and 4-(2'-Cyanomethylene-spiro[cyclohexane-1,3'-[3H]indol]-5'-yl)-thiophene-2-carbonitrile, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.

43(Withdrawn). The method according to claim 38, wherein said compound is 5'-(5-Cyano-1-methyl-1H-pyrrol-2-yl)spiro[cyclohexane-1,3'-[3H]indol]-2'-ylidenecyanamide, or a pharmaceutically acceptable salt, tautomer, metabolite, or prodrug thereof.